



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\sin\left(\phi + \frac{\theta}{2}\right) \cdot \left\{ \cos^2\left(\phi + \frac{\theta}{2}\right) + \sec^2 \frac{\theta}{2} \right\} = 0,$$

$$\text{whence } \phi = -\frac{\theta}{2}.$$

That is, the medial line is horizontal. The second factor gives imaginary results, except when  $\theta=0$ .

---

## PROBLEMS.

---

39. Proposed by B. F. FINKEL, A. M., Professor of Mathematics and Physics in Drury College, Springfield, Missouri.

A person whose height is  $a$  and weight  $W$  stands in a swing whose length is  $l$ . Supposing the initial inclination of the swing to the vertical is  $\alpha$  and that the person always crouches when in the highest position and stands up when in the lowest, his center of gravity moving through a distance  $b$  measured from lower part of swing upward, find how much the arc is increased after  $n$  complete vibrations.

40. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania.

Find the law of the force, in order that the orbit may be a Cassinian Oval.

41. Proposed by O. W. ANTHONY, M. Sc., Professor of Mathematics and Astronomy, New Windsor College, New Windsor, Maryland.

If the earth were a perfect sphere and had a frictionless surface, what would be the motion of a ball placed at a given latitude?

---

## DIOPHANTINE ANALYSIS.

---

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

---

## SOLUTIONS OF PROBLEMS.

---

40. Proposed by F. P. MATZ, D. Sc., Ph. D., Professor of Mathematics and Astronomy in Irving College, Mechanicsburg, Pennsylvania.

The sum of three positive integral *cubic* roots of an equation is a square. What is the equation?

I. Solution by E. L. SHERWOOD, A. M., Professor of Mathematics and Science in Mississippi Normal College, Houston, Mississippi.

Let  $a$ ,  $b$ , and  $c$  be the roots of the equation.

We then have  $a^3 + b^3 + c^3 = \square$ .

This condition is satisfied by the equation  $v^4(v^2 + 8v^2 + 27v^2) = \square$ , where